

**Recommended Adult\* Equine Preventative Vaccination Program** (10/09)  
 (\*Horses 1 year of Age or Older)

VACCINE	ADMINISTERED <i>IM - Intramuscular</i> <i>IN - Intranasal</i>	FREQUENCY
Tetanus/Sleeping Sickness <sup>1</sup>	IM	Annually <sup>4</sup>
Tetanus/Sleeping Sickness/West Nile Virus Combination	IM	Annually <sup>3 &amp; 4</sup>
Rabies	IM	Annually <sup>4</sup>
Flu/Rhino or Flu	IM or IN	Every 6 months
Strangles (IN) <sup>2</sup>	IN	Annually <sup>4</sup>
Prevenile (West Nile)	IM	Annually <sup>5</sup>

<sup>1</sup> In case of a penetrating wound, Tetanus Toxoid should be given if greater than 6 months since vaccination. If no prior Tetanus Toxoid can be established, consider Tetanus Antitoxin.

<sup>2</sup> We recommend the use of this vaccine only where there is known or possible exposure.

<sup>3</sup> If this vaccine is given in late March to a previously vaccinated horse, a West Nile booster in the same year should not be necessary

<sup>4</sup> Previously unvaccinated adults should be boosted 3-4 weeks after their initial vaccine.

<sup>5</sup> This West Nile vaccine is recommended for foals and other previously unvaccinated horses.