



Loomis Basin Equine Medical Center, Inc.

2973 Penryn Road Penryn, CA 95663
(916) 652-7645 Fax: (916) 652-7662 www.lbemc.com

Recommended Adult* Equine Preventative Vaccination Program ^(1/16)

(*Horses 1 year of Age or Older)

VACCINE	ADMINISTERED <i>IM - Intramuscular</i> <i>IN - Intranasal</i>	FREQUENCY
Tetanus/Sleeping Sickness ¹	IM	Annually ⁴
Tetanus/Sleeping Sickness/West Nile Virus Combination	IM	Annually ^{3 & 4}
Rabies	IM	Annually ⁴
Flu/Rhino or Flu	IM or IN	Every 6 months
Strangles (IN) ²	IN	Annually ⁴
West Nile	IM	Annually ^{4 & 5}

¹ In case of a penetrating wound, Tetanus Toxoid should be given if greater than 6 months since vaccination. If no prior Tetanus Toxoid can be verified, consider using Tetanus Antitoxin.

² We recommend the use of this vaccine only where there is known or possible exposure.

³ If this vaccine is given in late March to a previously vaccinated horse, a West Nile booster in the same year should not be necessary

⁴ Previously unvaccinated adults should be boosted 3-4 weeks after their initial vaccine.

⁵ This vaccine should be given in March or April (once yearly) if the Tetanus/Sleeping Sickness/West Nile combination is not used.